



Survival exponents for fractional Brownian motion with multivariate time

George Molchan

Institute of Earthquake Prediction Theory and Mathematical Geophysics,
Russian Academy of Science,
84/32 Profsoyuznaya st., 117997, Moscow, Russian Federation.
E-mail address: molchan@mitp.ru

Abstract. Fractional Brownian motion of index $0 < H < 1$, H -FBM, with d -dimensional time is considered in a spherical domain that contains 0 at its boundary. The main result: the log-asymptotics of the probability that H -FBM does not exceed a fixed positive level is $(H - d + o(1)) \log T$, where $T \gg 1$ is the radius of the domain.

1. Introduction

Fractional Brownian motion of index $H \in (0, 1)$, H -FBM, with multivariate time $t \in R^d$ is a centered Gaussian random process $w_H(t)$ with correlation function

$$E w_H(t) w_H(s) = 0.5(|t|^{2H} + |s|^{2H} - |t - s|^{2H}).$$

H -FBM is H -self-similar (H -ss), isotropic, and has stationary increments (si), i.e.,

$$\{w_H(\lambda U t + t_0) - w_H(t_0)\} \doteq \{\lambda^H w_H(t)\}$$

holds in the sense of the equality of finite-dimensional distributions for any fixed t_0 , $\lambda > 0$, and orthogonal mapping $U : R^d \rightarrow R^d$.

The one-sided exit problem for a random process $\xi(t)$ and its characteristics, the so-called survival exponents:

$$\theta_\xi = \lim_{T \rightarrow \infty} -\log P(\xi(t) < 1, t \in \Delta_T) / \psi(T) \quad (1.1)$$

are the subject of intensive analysis in applications. Here Δ_T is an increasing sequence of domains of size T , and $\psi(\cdot)$ is a suitable slowly varying function, typically, $\psi(T) = \log T$ for ss-processes. The greatest progress in this area has been achieved for processes with one-dimensional time. (See surveys by [Bray et al., 2013](#) of the

Received by the editors September 27, 2016; accepted December 2, 2016.

2010 *Mathematics Subject Classification.* 60G22.

Key words and phrases. Fractional Brownian motion; One-sided exit problem; Survival exponent.

Research supported by Russian Science Foundation, grant 17-11-01052.

physics literature and by [Aurzada and Simon, 2015](#) of the relevant mathematical publications).

H-FBM was one of the first non-trivial examples of non-Markovian processes for which the survival exponents have been found exactly ([Molchan, 1999](#)). Namely, for $\psi(T) = \log T$, the survival exponents for H-FBM are

$$\theta_{w_H} = 1 - H, \quad \Delta_T = (0, T) \quad \text{and} \quad \theta_{w_H} = d, \quad \Delta_T = (-T, T)^d. \quad (1.2)$$

Recently, [Aurzada et al. \(2016\)](#) considerably refined the asymptotics of probability

$$p_T = P(w_H(t) < 1, t \in \Delta_T), \quad \Delta_T = (0, T) \quad (1.3)$$

and showed that the exponent $\theta = 1 - H$ is universal for a broad class of H-ss processes with stationary increments. The ideas of this work have proved useful in the analysis of the conjecture that $\theta_{w_H} = d - kH$ for $w_H(t)$ in $\Delta_T = [0, T]^k \times [-T, T]^{d-k}$ ([Molchan, 2012](#)).

The case $k = 0$ corresponds to the right part of (1.2). The case $k = 1$ is supported by the result which we discuss below: $\theta_{w_H} = d - H$ for fractional Brownian motion in $\Delta_T = T\Delta_1$, where Δ_1 is a unit ball that contains 0 at its boundary.

The main idea of the paper by [Aurzada et al. \(2016\)](#) is to show that for a broad class of si-processes, $\xi(t), \xi(0) = 0$, with discrete time

$$|\Delta_T| P(\xi(t) < 1, t \in \Delta_T \cap \mathbb{Z}^1) \approx E \max(\xi(t), t \in \Delta_T \cap \mathbb{Z}^1), \quad (1.4)$$

where $\Delta_T = [0, T]$, $|\Delta_T| = T$, and \approx means up to a multiplicative term in $T^{o(1)}$. For H-ss processes with continuous time, the right-hand part of (1.4) is proportional to T^H , and therefore the exponent for (1.3) is $1 - H$. However, the result by [Aurzada et al. \(2016\)](#) essentially uses the 1-D nature of time. Considering $|\Delta_T|$ as the volume of Δ_T , relation (1.4) is found to be in formal agreement with the conjecture for $k = 1$, but not for $k > 1$; in addition, (1.4) is very crude for $k = 0$ (see (1.3)). This means that the analysis of the cases $d > 1, k > 1$ needs more ideas.

2. The lower bound

Proposition 2.1. *Let $\xi(t), \xi(0) = 0, t \in \mathbb{R}^d$ be a centered isotropic random process with stationary increments. Then*

$$P(\xi(t) < -1, t \in \Delta_T, |t| > 1) \leq cT^{-d} E \max(\xi(t), t \in \Delta_T),$$

where $\Delta_T = T\Delta_1$ is a ball of radius T that contains 0 at its boundary.

Consequence 2.2. If $\xi(t)$ is fractional Brownian motion of index $H \in (0, 1)$ in Δ_T , then the survival exponent has the lower bound $\theta_{w_H}^- \geq d - H$.

Remark 2.3. Proposition 2.1 holds for $\Delta_T = [0, T] \times [-T, T]^{d-1}$ as well.

Proof: Let $U_T = \{x_{k,\alpha}, \alpha = 1, 2, \dots, n_k; k = 1, 2, \dots\}$ be a subset of ball B_T of radius T in \mathbb{R}^d ; U_T consists of N_T points such that

$$|x_{k,\alpha}| = r_k, \quad |x_{k,\alpha} - x_{m,\beta}| > 1, \quad N_T > CT^d, \quad 1 < r_k < r_{k+1} \leq T. \quad (2.1)$$

Consider the following increasing sequence of subsets of U_T :

$$U_{k+1,\alpha} = U_k \cup \bigcup_{\beta=1}^{\alpha} x_{k+1,\beta}, \quad U_k = \{x_{i,\gamma} : |x_{i,\gamma}| \leq r_k\}.$$

Fix $\Delta_T = \{t : |t + Te| \leq T\}$, where $e = (0, \dots, 0, 1)$. Let $O_{k,\alpha}$ be an orthogonal mapping transforming $x_{k,\alpha}$ in $\tilde{x}_{k,\alpha} = r_k e$. Setting $\tilde{U}_{k,\alpha} = O_{k,\alpha} U_{k,\alpha}$, one has

$$(\tilde{U}_{k,\alpha} - \tilde{x}_{k,\alpha}) \setminus \{0\} \subset \Delta_T \setminus B_1, \quad (k, \alpha) \neq (1, 1). \quad (2.2)$$

Therefore, using the notation $M(A) = \sup(\xi(t), t \in A)$, we get

$$p_T(-1) := P(\xi(t) < -1, t \in \Delta_T \setminus B_1) \leq P(M((\tilde{U}_{k,\alpha} - \tilde{x}_{k,\alpha}) \setminus \{0\}) < -1). \quad (2.3)$$

By the si-property of $\xi(t)$, we can continue

$$= P(M(\tilde{U}_{k,\alpha} \setminus \tilde{x}_{k,\alpha}) - \xi(\tilde{x}_{k,\alpha}) < -1) = P(M(U_{k,\alpha-1}) + 1 < \xi(x_{k,\alpha})). \quad (2.4)$$

The last equality holds because $\xi(t)$ is rotation invariant.

The event $\{M(U_{k,\alpha-1}) + 1 < \xi(x_{k,\alpha})\}$ is measurable relative to the sequence

$$\xi(x_{1,1}), \dots, \xi(x_{1,n_1}); \dots; \xi(x_{k,1}), \dots, \xi(x_{k,n_k}); \dots \quad (2.5)$$

This event take place when $\xi(x_{k,\alpha})$ is realized as a record in the sequence (2.5) which exceeds the previous one by at least 1. Let ν_T be the number of such records in (2.5). Then, by (2.3, 2.4),

$$(N_T - 1)p_T(-1) \leq \sum_{k,\alpha} P(M(U_{k,\alpha}) + 1 < \xi(x_{k,\alpha+1})) = E\nu_T \leq E(M(U_T) - \xi(x_{1,1})),$$

where $U_{1,1} = \{x_{1,1}\}$, $U_{k,n_k} = U_{k+1}$, $x_{k,n_k+1} = x_{k+1,1}$, $(k, \alpha) \neq (1, 1)$.

Finally, by (2.1),

$$p_T(-1) \leq E(M(U_T)) / (N_T - 1) < cT^{-d} E(\sup \xi(t), t \in \Delta_T). \quad (2.6)$$

Suppose that $\xi(t)$ is fractional Brownian motion of index $H \in (0, 1)$ in Δ_T . By the standard procedure, we can compare $p_T(-1)$ with

$$p_T(1) = P(w_H(t) < 1, t \in (\Delta_T \setminus B_1)). \quad (2.7)$$

For this purpose we can find a continuous function $\varphi_T(t)$ such that

$$\varphi_T(t) = 1, |t| > 1, \|\varphi_T\|_{H,T} < const, \quad (2.8)$$

where $\|\cdot\|_{H,T}$ is the norm of the Hilbert space $H_H(\Delta_T)$ with the reproducing kernel $Ew_H(t)w_H(s)$, $(t, s) \in \Delta_T \times \Delta_T$ (see for this fact [Molchan, 1999](#) or Appendix). Then

$$p_T(-1) = P(w_H(t) + 2\varphi_T(t) < 1, t \in (\Delta_T \setminus B_1)).$$

According to [Aurzada and Dereich \(2013\)](#),

$$\left| \sqrt{\ln 1/p_T(1)} - \sqrt{\ln 1/p_T(-1)} \right| \leq \|2\varphi_T\|_{H,T} / \sqrt{2}. \quad (2.9)$$

From the self-similarity of H-FBM and (2.6) one has

$$p_T(-1) \leq cT^{-(d-H)} EM_{w_H}(\Delta_1). \quad (2.10)$$

Combining (2.8-2.10), one has

$$[\ln 1/P(w_H(t) < 1, t \in \Delta_T)]^{1/2} / \sqrt{\ln T} \geq \sqrt{d-H} + O(1/\sqrt{\ln T}), \quad (2.11)$$

i.e., $\theta_{w_H} \geq d - H$. \square

3. The upper bound

Below we use notation $M(A) = \sup(w_H(t), t \in A)$ and $|A| = \#\{t : t \in A\}$.

Proposition 3.1. *Let $w_H(t)$ be H-FBM in $\Delta_T = T\Delta_1 \subset R^d$ where Δ_1 is a bounded domain and $0 \in \Delta_1$. Consider a finite 1-net of Δ_T , i.e. a subset $U_T = \{x_k, k = 1, \dots, N_T\} \subset \Delta_T$, $\{0\} \notin U_T$ such that*

$$c < N_T/T^d < C \quad \text{and} \quad \Delta_T \subset \bigcup_{r=1}^{N_T} B_1(x_r),$$

where $B_1(x)$ is a unit ball centered at x . Then there is a $0 < q < 1$ such that for all $T > T_0$

$$P(M(\Delta_T) < c_H \sqrt{4d \ln T}) \geq qP(M(U_T) < 0). \quad (3.1)$$

In addition,

$$EM(U_T) = EM(\Delta_T)(1 + o(1)) = T^H EM(\Delta_1)(1 + o(1)), \quad T \rightarrow \infty. \quad (3.2)$$

Proof: One has

$$P(M(U_T) < 0) \leq P(M(U_T) < 0, A_T) + P(A_T^c), \quad (3.3)$$

where

$$A_T = \{\max_k \max_t (w_H(t) - w_H(x_k), t \in B_1(x_k)) < b_T\}.$$

We can continue the previous inequality

$$\begin{aligned} &\leq P(M(\Delta_T) < b_T) + \sum_k P(\max_t (w_H(t) - w_H(x_k), t \in B_1(x_k)) > b_T) \\ &\leq P(M(\Delta_T) < b_T) + N_T P(M(B_1) > b_T) := p_{1,T} + p_{2,T}. \end{aligned} \quad (3.4)$$

Applying the [Fernique \(1975\)](#) inequality to $w_H(t)$, we have

$$P(M(B_1) > r_T c_H) \leq c_d \int_{r_T}^{\infty} e^{-u^2/2} du, \quad r_T > (1 + 4d)^{1/2}. \quad (3.5)$$

Hence, setting $b_T = \sqrt{2(2d + \varepsilon) \ln T} c_H$, $\varepsilon > 0$, one has

$$p_{2,T} < CT^d \cdot T^{-2d - \varepsilon} / \sqrt{\ln T} = CT^{-d - \varepsilon} / \sqrt{\ln T}. \quad (3.6)$$

To show that $p_{2,T} = o(p_{1,T})$, note that $\Delta_T \subset B_{TD}$, where D is the diameter of Δ_1 . Therefore

$$p_{1,T} = P(M(\Delta_T) < b_T) \geq P(M(B_{TD}) < b_T) = P(M(B_{T'}) < 1), \quad (3.7)$$

where $T' = TD/b_T^{1/H}$. By [Molchan \(1999\)](#),

$$P(M(B_{T'}) < 1) > cT^{-d - \varepsilon}.$$

Due to (3.6), (3.7), we have

$$p_{2,T}/p_{1,T} < c(\ln T)^{-(1+d/H)/2} = o(1). \quad (3.8)$$

Relations (3.3), (3.4) and (3.8) imply (3.1):

$$P(M(U_T) < 0) \leq (1 + o(1))p_{1,T} \leq (1 + \varepsilon)P(M(\Delta_T) < b_T),$$

where $b_T = \sqrt{4d \ln T} c_H$. To prove relation (3.2), note that

$$\begin{aligned} M(\Delta_T) &\leq M(U_T) + \max_k \max_t (w_H(t) - w_H(x_k), t \in B_1(x_k)) \\ &:= M(U_T) + \delta_T. \end{aligned}$$

As above, using the event $A_T = \{\max_k \max_t (w_H(t) - w_H(x_k), t \in B_1(x_k)) < b_T\}$, one has

$$E\delta_T \leq b_T + E\delta_T 1_{A_T^c} \leq b_T + N_T EM(B_1)[M(B_1) > b_T], \quad (3.9)$$

where $b_T = \sqrt{4d \ln T} c_H$ and $N_T < CT^d$.

Therefore, the second term in (3.9) is $o(1)$, because

$$(EM(B_1)[M(B_1) > b_T])^2 \leq EM^2(B_1)P(M(B_1) > b_T) = O(T^{-2d}/\sqrt{\ln T}).$$

Due to (3.9), the relation (3.2) follows from the inequality:

$$\begin{aligned} EM(U_T) &\geq EM(\Delta_T) - E\delta_T \geq EM(\Delta_T) - c\sqrt{\ln T} + o(1) \\ &= T^H EM(\Delta_1) - c\sqrt{\ln T} + o(1). \end{aligned}$$

□

Proposition 3.2. *Let $w_H(t)$, $t \in \Delta_T$ be H-FBM, $\Delta_T = T\Delta_1 \subset R^d$, where Δ_1 is a unit ball and $0 \in \Delta_1$. Then*

$$P(M(\Delta_T) < 1) \geq cT^{-(d-H)}(\sqrt{\ln T})^{-d/H},$$

i.e., the survival exponent for H-FBM in Δ_T has the upper bound $\theta_{w_H}^+ \leq d - H$.

Corollary 3.3. *Due to Propositions 2.1, 3.2, the survival exponent for H-FBM in Δ_T exists and equals $d - H$.*

Proof: Proceeding as in the proof of Proposition 2.1, we consider again the subset U_T of the ball $B_T \subset R^d : U_T = \{x_{k,\alpha}, \alpha = 1, 2, \dots, n_k; k = 1, 2, \dots\}$, $\{0\} \notin U_T$. In addition to the properties (2.1), we suppose that the elements of U_T are enumerated in such a way that

$$x_{k,\alpha+1} \in B_2(x_{k,\alpha}) \quad \text{and} \quad x_{k+1,1} \in B_2(x_{k,n(k)}). \quad (3.10)$$

As before,

$$U_{k+1,\alpha} = U_k \cup \bigcup_{\beta=1}^{\alpha} x_{k+1,\beta}, \quad U_k = \{x_{i,\gamma} : |x_{i,\gamma}| \leq r_k\} := U_{k,0};$$

$\Delta_T = \{t : |t + Te| \leq T\}$, where $e = (0, \dots, 0, 1)$; $O_{k,\alpha}$ is an orthogonal mapping transforming $x_{k,\alpha}$ in $\tilde{x}_{k,\alpha} = r_k e$. Setting $\tilde{U}_{k,\alpha} = O_{k,\alpha} U_{k,\alpha}$, one has

$$(\tilde{U}_{k+1,\alpha} - \tilde{x}_{k+1,\alpha}) \setminus \{0\} \subset \Delta_{k+1} \setminus B_1.$$

Due to (3.10), $(\tilde{U}_{k+1,\alpha} - \tilde{x}_{k+1,\alpha})$ is a 2-net in Δ_{k+1} . Therefore, by (3.1), for $k > T_0$

$$\begin{aligned} P\left(M(\Delta_k) < c_H \sqrt{4d \ln k}\right) &> qP(M(\tilde{U}_{k,\alpha} - \tilde{x}_{k,\alpha}) \setminus \{0\} < 0) \\ &= qP(M(\tilde{U}_{k,\alpha-1}) - w_H(\tilde{x}_{k,\alpha}) < 0) = qP(M(U_{k,\alpha-1}) < w_H(x_{k,\alpha})). \end{aligned}$$

As a result,

$$\sum_{k=K}^{K'} n_k P\left(M(\Delta_k) < c_H \sqrt{4d \ln k}\right) \geq q \sum_{k=K}^{K'} \sum_{\alpha=1}^{n_k} P(M(U_{k,\alpha-1}) < w_H(x_{k,\alpha})) \quad (3.11)$$

where $K = [T]$ and $K' = [T']$.

Similarly to the proof of Proposition 2.1, we conclude that the right-hand part of (3.11) is equal to $qE\nu(T, T')$, where $\nu(T, T')$ is the number of all records in the following sequences:

$$M(U_K), w_H(x_{K+1,1}), \dots, w_H(x_{K+1,n(K+1)}); \dots; w_H(x_{K',1}), \dots, w_H(x_{K',n(K')}).$$

Let $\delta(T, T')$ be the maximum increment between adjacent elements of the sequence $w_H(x_{K,n(K)}), w_H(x_{K+1,1}), \dots, w_H(x_{K+1,n(K+1)}); \dots; w_H(x_{K',1}), \dots, w_H(x_{K',n(K')})$. Then

$$M(U_{K'}) - M(U_K) \leq (\nu(T, T') + 1)\delta(T, T') \leq (\nu(T, T') + 1)b_T + R_T, \quad (3.12)$$

where

$$R_T = (|U_{K'} \setminus U_K| + 1)\delta(T, T')[\delta(T, T') > b_T].$$

Due to (3.10),

$$ER_T < (|U_{K'} \setminus U_K| + 1)^2 \max_{|t| < 2} Ew_H(t)[w_H(t) > b_T].$$

Setting $b_T = \sqrt{8d \ln T} c_H$ and $T' - T = \rho T$, we obtain

$$ER_T < cT^{2d} \cdot T^{-2d} = c. \quad (3.13)$$

By (3.12),

$$b_T E\nu(T, T') > EM(U_{K'}) - EM(U_K) - b_T - ER_T,$$

where, according to (3.2),

$$EM(U_K) = K^H EM(\Delta_1)(1 + o(1)).$$

As a result,

$$b_T E\nu(T, T') > c(T^H - \sqrt{\ln T} - 1) = cT^H(1 + o(1)). \quad (3.14)$$

Keeping in mind that the right part of (3.11) is $qE\nu(T, T')$, we have:

$$qE\nu(T', T) \leq \sum_{k=K}^{K'} n_k P\left(M(\Delta_k) < c_H \sqrt{4d \ln k}\right). \quad (3.15)$$

Due to the self-similarity of H-FBM,

$$P\left(M(\Delta_k) < c\sqrt{\ln k}\right) = P\left(M(\Delta_{\tilde{k}}) < 1\right), \quad \tilde{k} = k / \left(c\sqrt{\ln k}\right)^{1/H},$$

and therefore the probability term decreases as a function of k . Hence, (3.15) implies

$$\begin{aligned} qE\nu(T', T) &\leq |U_{T'} \setminus U_T| P\left(M(\Delta_{T'+1}) < c_H \sqrt{4d \ln(T'+1)}\right) \\ &\leq CT^d P(M(\Delta_{\tilde{T}}) < 1), \end{aligned} \quad (3.16)$$

where

$$\tilde{T} = T' / \left(c_H \sqrt{4d \ln T'}\right)^{1/H} \quad \text{or} \quad T' = \tilde{T} \left(c_H \sqrt{4d \ln \tilde{T}}\right)^{1/H} (1 + o(1)). \quad (3.17)$$

Finally, by (3.14, 3.16),

$$b_T^{-1} cT^H (1 + o(1)) \leq E\nu(T', T) \leq q^{-1} CT^d P(M(\Delta_{\tilde{T}}) < 1).$$

Taking into account (3.17) and the relation $T' - T = \rho T$, we get

$$P(M(\Delta_{\tilde{T}}) < 1) \geq c\tilde{T}^{-(d-H)} \left(\sqrt{\ln \tilde{T}}\right)^{-d/H}.$$

□

Appendix

Example from Proposition 2.1. Consider H-FBM in domains $\Delta_T = T \cdot \Delta_1$, $0 \in \partial\Delta_1$; then a suitable function $\varphi_T(t)$, $t \in \Delta_T$ can be chosen as follows:

$$\varphi_T(t) = f(|t|/(Tk)) - f(|t|),$$

where $f(x)$, $x \in R^1$ is a finite smooth function such that $f(t) = 1$ for $|x| < 1/2$ and $f(t) = 0$ for $|x| > 1$. Here k is the diameter of Δ_1 .

By Molchan (1999), this can be seen as follows. Due to the spectral representation of H-FBM, the Hilbert space $H_H(\Delta_T)$ with the reproducing kernel $Ew_H(t)w_H(s)$, $(t, s) \in \Delta_T \times \Delta_T$ (Lifshits, 2012), is the closure of smooth functions $\varphi(t)$, $\varphi(0) = 0$ relative to the norm

$$\|\varphi\|_{H,T} = \inf_{\tilde{\varphi}} \|\tilde{\varphi}\|_H, \|\psi\|_H = A_H \int \left| \hat{\psi}(\lambda) \right|^2 |\lambda|^{d+2H} d\lambda,$$

Where $\tilde{\varphi}(t)$ is a finite function such that $\tilde{\varphi}(t) = \varphi(t)$, $t \in \Delta_T$; $\hat{\psi}(\lambda)$, $\lambda \in R^d$ is the Fourier transform of $\psi(t)$. Obviously, we have $\varphi_T(0) = 0$, $\varphi_T(1) = 1$ for $t \in \Delta_T \setminus B_1$, and

$$\begin{aligned} \|\varphi\|_{H,T} &< \|f(|t|/Tk) - f(|t|)\|_H < \|f(|t|/Tk)\|_H + \|f\|_H \\ &= ((Tk)^{-H} + 1) \|f\|_H < 2 \|f\|_H. \end{aligned}$$

Acknowledgements

I am very grateful to reviewer for careful reading of the paper and constructive comments.

References

- F. Aurzada and S. Dereich. Universality of the asymptotics of the one-sided exit problem for integrated processes. *Ann. Inst. Henri Poincaré Probab. Stat.* **49** (1), 236–251 (2013). [MR3060155](#).
- F. Aurzada, N. Guillotin-Plantard and F. Pene. Persistence probabilities for stationary increment processes. *ArXiv Mathematics e-prints* (2016). [arXiv: 1606.00236](#).
- F. Aurzada and T. Simon. Persistence probabilities and exponents. In *Lévy matters. V*, volume 2149 of *Lecture Notes in Math.*, pages 183–224. Springer, Cham (2015). [MR3468226](#).
- A. J. Bray, S. N. Majumdar and G. Schehr. Persistence and first-passage properties in nonequilibrium systems. *Advances in Physics* **62** (3), 225–361 (2013). [DOI: 10.1080/00018732.2013.803819](#).
- X. Fernique. Régularité des trajectoires des fonctions aléatoires gaussiennes. In *École d'Été de Probabilités de Saint-Flour, IV-1974*, pages 1–96. Lecture Notes in Math., Vol. 480. Springer, Berlin (1975). [MR0413238](#).
- M. Lifshits. *Lectures on Gaussian processes*. Springer Briefs in Mathematics. Springer, Heidelberg (2012). ISBN 978-3-642-24938-9; 978-3-642-24939-6. [MR3024389](#).
- G. Molchan. Maximum of a fractional Brownian motion: probabilities of small values. *Comm. Math. Phys.* **205** (1), 97–111 (1999). [MR1706900](#).
- G. Molchan. Survival exponents for some Gaussian processes. *Int. J. Stoch. Anal.* pages Art. ID 137271, 20 (2012). [MR2999109](#).